

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 31/70, C07H 15/12, 17/00 C12N 7/04	A1	(11) International Publication Number: WO 91/14436 (43) International Publication Date: 3 October 1991 (03.10.91)
(21) International Application Number: PCT/US91/01822 (22) International Filing Date: 19 March 1991 (19.03.91) (30) Priority data: 497,090 21 March 1990 (21.03.90) US (60) Parent Application or Grant (63) Related by Continuation US 497,090 (CIP) Filed on 21 March 1990 (21.03.90) (71) Applicant (for all designated States except US): ISIS PHARMACEUTICALS, INC. [US/US]; 2280 Faraday Avenue, Carlsbad, CA 92008 (US).		(72) Inventors; and (75) Inventors/Applicants (for US only) : ECKER, David, J. [US/US]; 2609 Colibri Lane, Carlsbad, CA 92009 (US). BRUICE, Thomas, W. [US/US]; 6880 Shearwaters Drive, Carlsbad, CA 92008 (US). VICKERS, Timothy, A. [US/US]; 907 Marlin Drive, Vista, CA 92084 (US). (74) Agents: CALDWELL, John, W. et al.; Woodcock Wash- burn Kurtz Mackiewicz & Norris, One Liberty Place, 46th Floor, Philadelphia, PA 19103 (US). (81) Designated States: AT (European patent), AU, BE (Euro- pean patent), BR, CA, CH (European patent), DE (Eu- ropean patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European pa- tent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, SE (European patent), US. Published <i>With international search report.</i>
(54) Title: REAGENTS AND METHODS FOR MODULATING GENE EXPRESSION THROUGH RNA MIMICRY (57) Abstract <p>Expression of genes may be modulated by employment of compositions which are capable of RNA mimicry. A portion of RNA coded by the gene whose expression is to be modulated is selected which is capable of interacting with one or more proteins. An oligonucleotide or oligonucleotide analog is then prepared in such a way as to mimic the portion of the RNA. Cells containing the gene are then contacted with the oligonucleotide or oligonucleotide analog to effect the modulation. Therapeutic compositions and methods, especially for the treatment of human immunodeficiency, are disclosed.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

-1-

**REAGENTS AND METHODS FOR MODULATING GENE
EXPRESSION THROUGH RNA MIMICRY**

FIELD OF THE INVENTION

This invention relates to the field of therapeutics,
5 particularly infections, in animals and humans. It relates
to the design, synthesis and application of oligonucleotide
analogs which mimic the RNA secondary structures found in
diseased cells, particularly cells infected with viruses and
retroviruses. These mimics of the infectious RNA structures
10 have been found to be able to modulate such infections.

BACKGROUND OF THE INVENTION

The biological function of RNA is mediated by its
structure. mRNA is generally thought of as a linear molecule
which contains the information for directing protein synthesis
15 within the sequence of ribonucleotides. Recently, studies
have revealed a number of secondary and tertiary structures
in mRNA which are important for its function (See; I. Tinoco,
P.W. Davis, C.C. Hardin, J.D. Puglisi, G.T. Walker, *Cold.
Spring. Harb. Symp. Quant. Biol.* 52, 135 (1987). Secondary
20 structural elements in RNA are formed largely by Watson-Crick
type interactions between different regions of the same RNA
molecule. Important secondary structural elements include
intramolecular double stranded regions, hairpin loops, bulges
in duplex RNA and internal loops. Tertiary structural
25 elements are formed when secondary structural elements come
in contact with each other or with single stranded regions to
produce a more complex, three dimensional structure.

Very little is known about the precise three

-2-

dimensional structure of RNA. However, there have recently been a number of research efforts which have shown that RNA structures, including single stranded, secondary, and tertiary structures, have important biological functions beyond simply encoding information to make proteins in linear sequences. Some of these correlations have been discussed in: I. Tinoco, P.W. Davis, C.C. Hardin, J.D. Puglisi, G.T. Walker, *Cold. Spring. Harb. Symp. Quant. Biol.* 52, 135 (1987); O. Resnekov, M. Kessler, Y. Aloni, *J. Biol. Chem.* 264, 9953 (1989); C. Tuerk, P. Gauss, C. Thermes, et al, *Proc. Natl. Acad. Sci. U. S. A.* 85, 1364 (1988); D.E. Larson, B.H. Sells, *Mol. Cell. Biochem.* 74, 5 (1987); and G. Knapp, *Methods Enzymol.* 180, 192 (1989).

Oligonucleotides have been evaluated for effect on HIV. Agarwal and coworkers have used oligonucleotide analogs targeted to the splice donor/acceptor site to inhibit HIV infection in early infected and chronically infected cells. S. Agarwal, T. Ikeuchi, D. Sun, P.S. Sarin, A. Konopka, J. Maizel, *Proc. Natl. Acad. Sci. USA* 86:7790 (1989). Sarin and coworkers have also used chemically modified oligonucleotide analogs targeted to the cap and splice donor/acceptor sites. P.S. Sarin, S. Agarwal, M.P. Civerira, J. Goodchild, T. Ikeuchi, P.C. Zamecnik, *Proc. Natl. Acad. Sci. USA* 85:7448 (1988). Zaia and coworkers have also used an oligonucleotide analog targeted to a splice acceptor site to inhibit HIV. Zaia, J.A., J.J. Rossi, G.J. Murakawa, P.A. Spallone, D.A. Stephens, B.E. Kaplan, *J. Virol.* 62:3914 (1988). Matsukura and coworkers have synthesized oligonucleotide analogs targeted to the initiation of translation of the rev gene mRNA. M. Matsukura, K. Shinozuka, G. Zon, et al., *Proc Natl. Acad. Sci. USA*, 84:7706 (1987); R.L. Letsinger, G.R. Zhang, D.K. Sun, T. Ikeuchi, P.S. Sarin, *Proc. Natl. Acad. Sci. USA* 86:6553 (1989). Mori and coworkers have used a different oligonucleotide analog targeted to the same region as Matsukura et al., K. Mori, C. Boiziau, C. Cazenave et al., *Nucleic Acids Res.* 17:8207 (1989). Shibahara and coworkers

-3-

have used oligonucleotide analogs targeted to a splice acceptor site as well as the reverse transcriptase primer binding site. S. Shibahara, S. Mukai, H. Morisawa, H. Nakashima, S. Kobayashi, N. Yamamoto, *Nucl. Acids Res.* 17:239 (1989). Letsinger and coworkers have synthesized and tested oligonucleotide analogs with conjugated cholesterol targeted to a splice site. K. Mori, C. Boiziau, C. Cazenave, et al., *Nucleic Acids Res.* 17:8207 (1989). Stevenson and Iversen have conjugated polylysine to oligonucleotide analogs targeted to the splice donor and the 5'-end of the first exon of the tat gene. M. Stevenson, P.L. Iversen, *J. Gen. Virol.* 70:2673 (1989). Each of these publications have reported some degree of success in inhibiting some function of the HIV virus. While each of these references is distinct from the approach of the present invention, each supports the view that nucleotide therapeutics in HIV infection is rational and based upon sound scientific principles. In each of these references the approach has been to design antisense oligonucleotides complementary to some portion of the HIV mRNA. The present invention relates to oligonucleotides which mimic an RNA and bind to a protein, rather than oligonucleotides which bind to the HIV RNA.

Heretofore, there have been no suggestions in the art of methods or materials which could be useful for mimicking the secondary or tertiary structures of RNA in order to modulate the expression of genes or to treat disease. This is despite the long-felt need for methods of therapeutics and for methods of inhibiting gene expression which may be related to diseases or disease states in animals. Accordingly, there remains a long-felt need for therapeutic materials and methods, especially for viruses and retroviruses.

OBJECTS OF THE INVENTION

It is a principal object of the invention to provide compositions and therapies for human diseases, particularly viral and retroviral infections.

-4-

It is a further object of the invention to provide therapeutic compositions which mimic the structure of a natural RNA.

Yet another object of this invention is to modulate
5 gene expression in cells.

Yet another object of this invention is to provide therapies for human immunodeficiency virus infection.

These and other objects of this invention will become apparent from a review of the instant specification.

10 SUMMARY OF THE INVENTION

It has now been discovered that expression of genes may be modulated through the employment of compositions which are capable of RNA mimicry. The use of such RNA mimics can interfere with gene expression and, when that expression is
15 implicated in the etiology of disease, lead to methods of therapeutics. In accordance with this invention, it has now been found that certain portions of RNA coded by genomic material can have secondary and even tertiary structure which plays a significant role in gene expression. It has now been
20 found that the interaction of certain RNA's, especially messenger RNA's having secondary or tertiary structures, with proteins may be inhibited through the employment of oligonucleotides or oligonucleotide analogs which mimic at least a portion of the RNA. Such mimicry can interfere with
25 the protein-RNA interaction and, through such interference, interfere with gene expression and the maintenance of disease states.

In accordance with preferred embodiments of the present invention, methods for modulating expression of a gene
30 are provided comprising selecting a portion of RNA coded by the gene, which RNA is capable of interacting with one or more proteins. An oligonucleotide or oligonucleotide analog is then prepared in such a way as to mimic said portion of the RNA. Cells containing the gene are then contacted with the
35 oligonucleotide or oligonucleotide analog to effect such modulation of expression. It will generally be the case that

-5-

the gene is of an infectious organism, such as a virus or retrovirus. Preferably, the gene is from human immunodeficiency virus.

In accordance with other preferred embodiments, the protein is produced by a second portion of RNA coded by the infectious organism such as a virus or retrovirus. In such a case, the interaction of the protein with the RNA portion selected, if permitted to occur, would generally effect stimulation of expression of the gene such that inhibition of this interaction effects repression or modulation of gene expression.

It is preferred that the oligonucleotide or oligonucleotide analogs of the invention mimic at least about 6 nucleotide units of the selected RNA. It is still more preferred that from 8 to about 60 nucleotide units be mimicked. From about 10 to about 30 nucleotide units are presently believed to be most preferred. In accordance with other preferred embodiments, the degree of mimicry of the selected RNA is such as to permit the oligonucleotide or oligonucleotide analog to achieve at least a portion of the secondary structure of the RNA.

In accordance with other preferred embodiments of the present invention, the TAR region, the CAR region, or the GAG-POL region of human immunodeficiency virus messenger RNA is targeted for oligonucleotide mimicry. The oligonucleotide or oligonucleotide analog is selected to be sufficient in its degree of mimicry as to be effective in interfering with the interaction of protein with the selected messenger RNA portions. Thus, for example, if the selected messenger RNA portion is the TAR region of HIV, then the oligonucleotide or oligonucleotide analog is constructed so as to mimic the TAR region sufficiently such that tat protein coded by another portion of the HIV messenger RNA is effectively complexed with or bound to the mimicking molecule. Similar considerations attend the preparation of oligonucleotide and oligonucleotide analog

-6-

RNA mimics directed at the CAR and GAG-POL regions of HIV.
BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1A depicts the linear HIV-1 TAR element sequence. Underlined portions connote loops and bulges.

5 Figure 1B depicts a computer-predicted secondary structure of the HIV-1 TAR element.

Figure 2 sets forth the partial linear structure of the HIV-1 CAR RNA sequence corresponding to nucleotides 7357-7627.

10 Figure 3 shows a computer-predicted secondary structure of the HIV-1 CAR element

Figure 4 shows the structure of oligonucleotides identified as 1345, 1346 and 1347.

15 Figure 5 is a graph showing the inhibition of HIV LTR gene expression observed with oligonucleotides 1345, 1346, 1347 and 1348.

Figure 6 is a graph showing the inhibition of HIV LTR gene expression observed with oligonucleotides 1345, 1349 and a control.

20 Figure 7 shows the activity of oligonucleotides identified as 2306, 2848 and 2850 for inhibition of HIV LTR gene expression.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

In accordance with the present invention,
25 compositions which mimic the structure of biological RNA molecules of significant importance are provided. The present invention employs oligonucleotides and oligonucleotide analogs to mimic the structures of the biological RNA molecules. In the context of this
30 invention, the term "oligonucleotide" refers to a plurality of joined nucleotide units formed from naturally-occurring bases and cyclofuranosyl groups joined by native phosphodiester bonds. This term effectively refers to naturally-occurring species or synthetic species formed
35 from naturally-occurring subunits.

-7-

"Oligonucleotide analog," as that term is used in connection with this invention, refers to moieties which function similarly to oligonucleotides but which have non-naturally occurring portions. Thus, oligonucleotide
5 analogs may have altered sugar moieties or inter-sugar linkages. Exemplary among these are the phosphorothioate and other sulfur containing species which are known for use in the art. They may also comprise altered base units or other modifications consistent with the spirit of this
10 invention.

In accordance with certain preferred embodiments, at least some of the phosphodiester bonds of the oligonucleotide have been substituted with a structure which functions to enhance the ability of the compositions
15 to penetrate into the region of cells where the RNA whose activity is to be modulated is located. It is preferred that such linkages be sulfur-containing. It is presently preferred that such substitutions comprise phosphorothioate bonds. Others such as alkyl phosphothioate bonds, N-alkyl
20 phosphoramidates, phosphorodithioates, alkyl phosphonates, and short chain alkyl or cycloalkyl structures may also be useful. In accordance with other preferred embodiments, the phosphodiester bonds are substituted with structures which are, at once, substantially non-ionic and non-chiral.
25 Persons of ordinary skill in the art will be able to select other linkages for use in the practice of the invention.

It is generally preferred for use in some embodiments of this invention that the 2' position of the linking sugar moieties in at least some of the subunits of
30 the oligonucleotides or oligonucleotide analogs be substituted. Thus, 2' substituents such as OH, SH, F, OCH₃, OCN, OCH_nCH₃ where n is from 1 to about 10 and other substituents having similar properties may be useful in some embodiments.

35 Oligonucleotide analogs may also include species which include at least some modified base forms. Thus,

purines and pyrimidines other than those normally found in nature may be so employed. Similarly, modifications on the cyclofuranose portions of the nucleotide subunits may also occur as long as the essential tenets of this invention are
5 adhered to.

Such analogs are best described as being functionally interchangeable with natural oligonucleotides (or synthesized oligonucleotides along natural lines), but which have one or more differences from natural structure.
10 All such analogs are comprehended by this invention so long as they function effectively to mimic the structure of the desired RNA.

The oligonucleotides and oligonucleotide analogs in accordance with this invention preferably comprise from
15 about 3 to about 100 subunits. It is preferred that such oligonucleotides and analogs comprise greater than about 6 subunits with from about 8 to about 60 subunits being more preferred, and still more preferred to have from about 10 to about 30 subunits. As will be appreciated, a subunit is
20 a base and sugar combination suitably bound to adjacent subunits through phosphodiester or other bonds.

The oligonucleotides and oligonucleotide analogs of this invention can be used in diagnostics, therapeutics and as research reagents and kits. For therapeutic use,
25 the oligonucleotide or oligonucleotide analog is administered to an animal, especially a human, such as is suffering from a virus or retrovirus infection such as AIDS.

It is generally preferred to apply the
30 therapeutic agent in accordance with this invention internally such as orally, intravenously or intramuscularly. Other forms of administration, such as transdermally, topically or intralesionally may also be useful. Inclusion in suppositories may also be useful.
35 Use of the oligonucleotides and oligonucleotide analogs of this invention in prophylaxis is also likely to be useful.

-9-

Use of pharmacologically acceptable carriers is also preferred for some embodiments. In accordance with the present invention, the oligonucleotides and oligonucleotide analogs which are useful in its performance are best described by the RNA which they are designed to mimic. Thus, it will be understood by persons of ordinary skill in the art that the oligonucleotides and analogs provided by this invention are those which are substantially identical to a portion of an RNA, especially a messenger RNA having a particular relationship to a diseased state. Thus, the RNAs which are to be mimicked in accordance with this invention are those RNAs having a secondary structure and which are capable of interacting with one or more proteins. While the present invention is not so limited, and while the inventors do not wish to be bound by any particular theory of operation of the present invention, it is believed that a number of regulatory centers are extant upon RNA coded by genes responsible for disease. It is believed that such regulatory RNA portions possess a secondary structure such as a hair pin loop, stem loop, bulge, or similar structure which is capable of interacting with a protein, generally a protein coded by a different portion of the same or different RNA. It is generally believed in some cases that the interaction of the protein or proteins with the regulatory RNA center causes or leads to an enhancement of translation of the RNA into protein. Overall, this is considered to be an enhancement in the expression of the underlying gene since, of course, the RNA is derived from said gene. The modulation of such enhancement is an object of this invention.

It has now been discovered that preparation of mimics to these regulatory RNA portions and placement of quantities of such mimics, which are oligonucleotides or oligonucleotide analogs, into the cells or tissues which are suffering from infection, can result in a diminution of the infection; a modulation of expression of the underlying

-10-

gene. This is believed to be effected by interaction of the protein or proteins with the mimic molecules such that interaction of the protein with the regulatory RNA portion is minimized. Accordingly, enhancement in the expression of the underlying gene is similarly modulated. The present invention is believed to be quite general in application. Thus, if an RNA is believed to be capable of interaction with a protein in a regulatory sense as discussed herein before, then design of an oligonucleotide or oligonucleotide analog which mimics the RNA or at least a portion thereof may lead to therapeutic materials and methods. Thus, by contacting an animal suffering from such infection with a mimicking oligonucleotide or oligonucleotide analog, diminution in the infection can be had.

While at present, the RNAs which are believed to have the regulatory relationship discussed above all appear to have secondary structures such as stem loops, bulges and the like, it is possible that regulatory RNA segments may exist which do not enjoy such secondary structures. In such case, the present invention is to be understood to contemplate the preparation of mimicking oligonucleotides or oligonucleotide analogs for such RNA as well. Similarly, this invention will be understood to extend to therapeutic methods and compositions for such RNAs.

A number of RNA secondary structures have recently been identified for which application of this invention will likely provide therapeutic utility. Some of these include the HIV TAR structures as reported by S. Feng, E.C. Holland, in *Nature* 334, 165 (1988); including the stem loops at nucleotide 5-54, and 58-104 according to the nucleotide sequence as described by Ratner in L. Ratner, W. Haseltine, R. Patarca, K.J. Livak, B. Starcich, S.F. Josephs, *Nature* 313, 277 (1985); the boundary between the EGP/OMP regions of HIV as disclosed by S. Le, J. Chen, M.J. Braun, M.A. Gonda, J.V. Maizel, in *Nucl. Acids Res.*

-11-

16, 5153 (1988); the boundary between the TMP/env genes of HIV (ibid), the HIV CAR structure as reported by E.T. Dayton, D.M. Powell, A.I. Dayton, in *Science* 246, 1625 (1989); and the stem loop structure at the junction between
5 the HIV gag and pol genes (nucleotides 1629-1674), the HIV CRS element, and the human iron responsive element (IRE) as described by J.L. Casey, M.W. Hentze, D.M. Koeller, et al., in *Science* 240, 924 (1988). In addition, there are regions of RNA which are primarily thought of as single stranded
10 areas which have been identified as sites for protein binding. For example, the sequence 5'-AUUUA-3' has been identified as a signal for a protein to bind which leads to degradation of RNA as disclosed by J.S. Malter, in *Science* 246, 664 (1989). The structure of this region is not
15 known. However, that does not preclude the practice of this invention with this sequence. Additional RNA elements, with as yet unknown structures, can also be the subject of this invention.

It is not necessary to know the actual RNA
20 structure in order to practice this invention, it is only necessary to know that a specific RNA sequence is recognized by an RNA binding protein and that this interaction has important biological consequences. In this regard, the viral RNA sequences and structures which are
25 recognized by the structural proteins of retroviruses for virion formation are also the subject of this invention. The mimicry of any RNA structure which may interact with protein to effect an important biological function may fall within the spirit and scope of this invention.

30 THE HIV TAR ELEMENT AND TRANSACTIVATION BY THE *tat* PROTEIN

The HIV TAR element provides a good example of this invention. An elaborate set of control elements in the HIV genome determines whether the virus replicates or remains dormant. Of the nine genes identified in the HIV
35 genome, only three are from the core and envelope as described by W.A. Haseltine, F. Wong-Staal, in *Scientific*

-12-

American, October, 52 (1988). The other six genes are involved in regulation of the production of viral proteins.

Regulatory genes work by encoding a protein that interacts with a responsive element somewhere else on the viral genome. The major regulatory gene responsible for initiating the burst of replication is the tat (trans-activator) gene. The product of the tat gene, tat protein, works by interaction with a short sequence element known as TAR (trans-acting responsive element). The TAR sequence is encoded in the viral long terminal repeats (LTR's), and therefore is included in the mRNA from every HIV gene.

Expression of the tat protein results in increased expression of other HIV genes up to 1,000 fold, including the tat gene itself. Because of this autoregulatory positive feedback, and the fact that the TAR sequence is included in the mRNA from every HIV transcript, an immense amount of viral gene expression is triggered when the tat gene is activated. The interaction between the tat gene and the TAR element is therefore crucial to the life cycle of HIV, and specific disruption of this interaction is believed likely to interrupt the propagation of the virus; to modulate gene expression.

The mechanism of trans-activation of TAR-containing genes by the tat protein has recently been studied intensely, as disclosed by Sharp, Philip A., and Marciniak, Robert, A., in Cell 59, 229 (1989). It was found that tat increases the expression of TAR-containing genes by increasing both the amount of viral mRNA and the efficiency of its translation. Moreover, it appears that TAR functions as an RNA structure, rather than a DNA structure. The surprising result is that tat increases the transcription of TAR-containing genes, but does so by interacting with the TAR element in RNA. In order to achieve trans-activation, the TAR element must be located immediately "downstream" from the site of initiation of

-13-

transcription. Moreover, TAR is orientation dependent; if inserted in the inverse orientation, it fails to function. TAR function does not depend upon the presence of other HIV sequences upstream of the initiation of transcription, but
5 will act independently of the promoter.

Some of the strongest evidence that tat interacts with TAR as an RNA structure has come from mutagenesis experiments. Efforts to study the TAR element and RNA structure were stimulated by the observation that the tat
10 protein from HIV-1 was capable of trans-activating vectors containing the TAR region of HIV-2, a different strain of virus, even though there is very little primary sequence homology in the TAR region between the two strains. See S. Feng, E.C. Holland, in *Nature* 334, 165 (1988). However,
15 examination of the TAR sequence from HIV-1 and HIV-2 with computer programs that predict RNA secondary structure revealed the potential of RNA stem-loop structures, with a single stem-loop in the TAR region of HIV-1 and three stem-loop structures in HIV-2. Although the compositions and
20 lengths of the stems were divergent, all four loops contained the pentanucleotide CUGGG. Figure 1 depicts the linear sequence of the HIV-1 TAR region with the feature underlined. Mutagenesis experiments by Feng, ibid, revealed that each of the nucleotides present in the loop
25 is essential for trans-activation by tat, but that base substitutions in the stem were tolerated so long as the stem structure was maintained. Figure 1A and 1B depict the linear (primary) and secondary structures of HIV-1 TAR.

Further evidence for the TAR structure function
30 was obtained from experiments in which the sequences flanking the stem-loop structure were altered creating competing secondary structures in the RNA that were more stable than the natural TAR stem-loop. See Ben Berkhout, in *Cell* 59, 273 (1989). This was accomplished by
35 introducing additional sequences into the TAR-containing RNA that were antisense to the 5' side of the stem-loop

-14-

structure. Trans-activation of the modified TAR structure was lost, suggesting that the TAR sequences alone are not sufficient for trans-activation, but that these sequences must fold up in the proper secondary structure to be
5 active. It also suggests that antisense sequences to the TAR stem-loop are capable of disrupting the natural RNA structure.

Direct biochemical evidence for TAR stem-loop structure has also been obtained. The TAR RNA has been
10 enzymatically synthesized *in vitro* and probed with enzymes which selectively cleave single stranded regions of RNA, but not duplex structures. The results of the cleavage patterns were consistent with the computer predicted RNA secondary structure.

15 Thus, it now appears that:

1. The HIV tat protein is responsible for triggering an enormous amount of viral gene expression;
2. This occurs by interaction with the TAR sequence which is incorporated into every HIV mRNA
20 transcript;
3. The HIV TAR sequence functions as an RNA secondary structure; and
4. The correct TAR RNA secondary structure is essential for tat transactivation.

25 Compounds have now been discovered which are believed to specifically mimic the TAR RNA structure and interfere with tat trans-activation. These oligonucleotide and oligonucleotide analog compounds will likely have activity as therapeutic agents for HIV infection.

30 It is intended that all strains of HIV fall within the spirit and scope of this invention. It will be realized that different strains of HIV will have different TAR sequences which will therefore fold into different structures. This invention can be practiced on alternative
35 strains of HIV by changing the sequence of the oligonucleotide or oligonucleotide analog to mimic the

-15-

structure of the alternative strain. Thus, this aspect of the invention relates to all such strains and to oligonucleotide mimics for each respective TAR region.

TAR and tat function has been studied by removing
5 the genes from the HIV genome and studying them in cell lines in isolation. Vectors have been constructed to study the interactions between the tat protein and TAR element. The tat gene is expressed under the SV40 promoter. The TAR region is expressed from a separate plasmid fused to an
10 easily assayed reporter gene such as the chloramphenicol acetyl transferase gene (CAT) or the placental alkaline phosphatase gene (PAP) as reported, for example, by S. Feng, E.C. Holland, in *Nature* 334, 165 (1988) and by P. Henthorn, P. Zervos, M. Raducha, H. Harris, T. Kadesch, in
15 *Proc. Natl. Acad. Sci. USA* 85, 6342 (1988).

Enzymatic activity in cell culture models has been shown to be dependent upon both the presence of the essential elements of the TAR region and the presence of the tat protein. Pertinent reviews include Philip A.
20 Sharp, Robert A. Marciniak, *Cell* 59, 229 (1989); Feng supra.; Michael F. Laspia, Andrew P. Rice, Michael B. Mathews, *Cell* 59, 283 (1989); J.A. Garcia, D. Harrich, E. Soultanakis, F. Wu, R. Mitsuyasu, R.B. Gaynor, *EMBO J.* 8, 765 (1989); and Ben Berkhout, *Cell* 59, 273 (1989). In
25 essence, the vector system reconstitutes the events of tat-mediated TAR transactivation which occurs in HIV infected cells.

tat/TAR transactivation can be assayed by placing the human placental alkaline phosphatase gene (PAP) under
30 the regulatory control of the HIV-1 LTR sequences, which contain enhancer, promoter, and TAR elements. A plasmid containing the HIV-1 LTR, pHIVCAT-0, as described by S. Feng, E.C. Holland, *Nature* 334, 165 (1988), contains HIV U3 in its entirety and R up through position +78 (a HindIII
35 site). Digestion of this plasmid with a combination of HindIII and AatII releases the CAT cassette along with the

-16-

SV40 sequences responsible for the processing of the RNA. A second plasmid, pSV2APAP, contains the PAP cassette with eukaryotic processing signals, under the transcriptional control of an SV40 promoter, as referenced by P. Henthorn, P. Zervos, M. Raducha, H. Harris, T. Kadesch, in *Proc. Natl. Acad. Sci. USA* 85, 6342 (1988). The PAP cassette and processing sequences can be released from the plasmid by digestion with HindIII and AatII. A new plasmid, pHIVPAP, was created by ligating the HindIII/AatII fragment containing the HIV-1 LTR and vector sequences from pHIVCAT-0, to the HindIII/AatII PAP cassette from pSV2APAP

It has been shown that pHIVCAT-0 is transactivated in the presence of a second plasmid, pcDEBtat, which expresses the tat coding region under the regulatory control of the SV40 promoter. However, no CAT activity is seen in the absence of co-transfection of pcDEBtat as disclosed by Feng. To test the activity of oligonucleotides and oligonucleotide analogs, pcDEBtat and pHIVPAP were co-transfected into HeLa cells using the calcium/phosphate method. 48 hours post-transfection cells were harvested and assayed for PAP activity as described by Henthorn et al. The effects of oligonucleotides and oligonucleotide analogs were determined by adding the compounds directly to the transfection mixture or by adding the compounds to the media at various times and concentrations following transfection, followed by PAP assay at, for example, 24-48 hours post-transfection.

Cells were treated with the following exemplary oligonucleotide and oligonucleotide analog sequences:

30 5'- -3'
 GGGUCUCUCUGGUUAGACCAGAUUCUGAGCCUGGGAGCUCUCUGGCUAACUAGGGGAACCC
 GGUUAGACCAGAUUCUGAGCCUGGGAGCUCUCUGGCUAACU
 UCUGAGCCUGGGAGCUCUCU
 CCAGAUUCUGAGCCUGGGAGCUCUCUGG
 35 GAGCCUGGGAGCUC
 CUGGGA

-17-

Modulation of HIV LTR gene expression, as monitored by PAP activity, was observed.

To be useful pharmacologically in the treatment of the previously described tat-mediated pathologies, TAR mimetics minimally must satisfy certain general structure/function criteria which are not adequately met by unmodified TAR RNA. The specific compositions of matter presented herein are designed to achieve the following goals. First and foremost, nuclease resistance (to RNases and RNA active DNases) must be conferred. Secondly, the minimal TAR fragment required for tat binding should be employed. Enhanced tat binding specificity and affinity, and therapeutic index, can be achieved by conformational stabilization of the preferred conformation of bound TAR. Finally, compositions could have enhanced affinity and specificity for tat by improvements on the natural chemical basis of specificity. A number of TAR mimetic oligonucleotide sequences have been prepared in accordance with the teachings of the invention as shown in Table 1:

20

TABLE 1

TAR MIMETIC OLIGONUCLEOTIDE SEQUENCES

<u>OLIGO</u>	<u>SEQUENCE</u>
TAR 59-mer 25 1-59	5'-GGG UCU CUC UGG UUA GAC CAG AUC UGA GCC UGG GAG CUC UCU GGC UAA CUA GGG AAC CC-3', SEQ. ID NO: 1
U-DNA-TAR #1973 40-mer 11-50	5'-GGU UAG ACC AGA UCU GAG CCU GGG AGC UCU CUG GCU AAC U-3', SEQ. ID NO: 2
30 A:P=S TAR 59-mer 1-59	5'-GGG UCU CUC UGG UUA GAC CAG AUC UGA GCC UGG GAG CUC UCU GGC UAA CUA GGG AAC CC-3', SEQ. ID NO: 1

-18-

loopless ΔTAR #2002/#2246 23-mer 16-29 (-A17) 5 +36-45	5'-GCC AGA UCU GAG C-3', SEQ. ID NO: 3
5-BrU TAR 59-mer 1-59	5'-GCU CUC UGG C-3', SEQ. ID NO. 4
10 2'-OMe ΔTAR #2306 29-mer 16-45(-A17)	5'-GGG UCU CUC UGG UUA GAC CAG AUC UGA GCC UGG GAG CUC UCU GGC UAA CUA GGG AAC CC-3', SEQ. ID NO: 1
15 2' OMe, P=S ΔTAR #2195 29-mer 16-45(-A17)	5'-GCC AGA UCU GAG CCU GGG AGC UCU CUG GC-3', SEQ. ID NO: 5
1345 18-mer 23-40	5'-GCC AGA UCU GAG CCU GGG AGC UCU CUG GC-3', SEQ. ID NO: 5
20 1346 58-mer 1-58	5'-UCU GAG CCU GGG AGC UCU-3'
25 1347 32-mer 23-54	5'-GGG UCU CUC UGG UUA GAC CAG AUC UGA GCC UGG G AGC UCU CUG GCU AAC UAG GGA ACC
1348 17-mer 38-54	5'-UCU GAG CCU GGG AGC UCU CUG GCU AAC UAG GG-3'
30 1349 25-mer 1-25	5'-UCU CUG GCU AAC UAG GG-3'
	5'-GGG UCU CUC UGG UUA GAC CAG AUC U-3'

We have shown that the compound 2306, a 2'-O-methyl oligonucleotide analog comprising the sequence shown in Table 1, has significant activity in inhibition of HIV gene expression.

Sullenger et al. disclose that expression of high levels of tRNA-TAR fusion transcripts correlated with effective inhibition of HIV-1 replication and prevented the cytopathic effects associated with HIV-1 replication in CEM SS cells. CEM SS is a human T-lymphoid cell line that is

-19-

highly susceptible to HIV-1 replication. Sullenger et al. believe it is reasonable to assume that tat must physically associate with TAR in order to assert its function, whether binding directly or indirectly via a cellular factor. If
5 so, overexpression of an RNA species encoding the TAR sequence could act as a decoy to bind tat and/or the cellular factor and prevent its binding to the TAR sequence encoded in the viral DNA. The result will be no activation of viral gene expression and no generation of progeny
10 virus. TAR decoy-mediated inhibition of HIV-1 replication in CEM SS cells is shown to be very efficient. Base changes in the TAR stem or loop sequence which abolish tat-mediated trans-activation are also shown to abolish the ability of TAR decoy RNA to inhibit HIV replication in
15 these cells. These results suggest but do not prove that HIV replication is inhibited in TAR decoy-containing cells because tat-mediated trans-activation is competitively squelched by the presence of an excess of nonviral TAR-containing RNA.

20 Graham et al., *Proc. Natl. Acad. Sci., USA* 87:5817-5821 (1990) disclose that a possible method of inhibiting tat-TAR interaction is to provide an excess of TAR decoys, i.e., TAR sequences (DNA or RNA) that competitively bind factors mediating transactivation and
25 prevent them from acting. A problem in the use of TAR decoys may be the inability to put enough copies into a target cell to be effective. A suggested solution to this problem is to assemble many copies of the TAR in a head-to-tail tandem array and insert them as a single
30 transcriptional unit, ideally behind a strong promoter. Graham et al. constructed an array of 12 TAR copies behind a strong promoter, the human β -actin promoter, and showed that the transcripts so produced in human cells do interfere with the tat-TAR interaction *in vivo*.

35 Although it has been shown that inhibition of HIV replication can be achieved without causing damage to

-20-

cells, no one, until now, has been able to make an RNA therapeutic compound which can inhibit viral replication *in vivo*. In the present invention, RNA mimetics, with modifications conferring stability, are employed to maintain the TAR decoy function. These RNA mimics are distinct from the TAR decoys described by the prior art, which introduce DNA into the cell which is then transcribed into RNA in the cell; resulting RNA (TAR) has not been useful therapeutically because of its instability.

10 tat AND KAPOSI'S SARCOMA

Kaposi's sarcoma (KS) is one of the diagnostic signs of AIDS, and is the initial manifestation in approximately 30% of patients with AIDS (Tom, *Hawaii Med. J.*, 48:131-134 (1989)). Skin lesions are the usual initial presentation, and can be single or multiple and range from faint pink to dark purple. The mucous membranes can be involved. In advanced disease, pain can be prominent; disfigurement and severe edema can also occur. Visceral involvement is also common in advanced disease; the tumors have been found in liver, spleen, gastrointestinal tract, oropharynx, conjunctiva, brain, testes, lungs, pancreas, aorta and heart. Kaposi's sarcomas are vascular tumors characterized by the proliferation of abnormal endothelial cells with spindle-shaped cells and extravasated red blood cells. Lesions in initial stages of disease are usually multifocal rather than metastatic. Metastases can occur, but usually late in the course of disease.

KS associated with AIDS differs from the previously known KS, which was rare and afflicted elderly men of Jewish or Mediterranean descent almost exclusively. This classic KS was usually indolent in its progression and required minimal treatment. In contrast, KS associated with AIDS is aggressive and is acknowledged as a significant cause of morbidity and mortality in AIDS patients. Treatment of AIDS-associated KS is largely experimental, nonspecific, and not very encouraging. Tom,

-21-

Hawaii Med. J., 48:131-134(1989) describes the disease in AIDS patients. The use of chemotherapy in these patients is controversial because it can further impair cellular immunity and increase the risk of opportunistic infections.

5 Trials with single agents such as the vinca alkaloids, vincristine and vinblastine, and a podophyllotoxin, VP-16, have shown variable results and mild-to-moderate toxicity. However, response duration is short, and relapses frequent. Combination therapy has also been tried but is associated
10 with a significant incidence of opportunistic infections. Immunotherapy has also been tried; alpha interferon has been shown to be active; however, this may be due to its antitumor effect. No treatment to date has resulted in any reversal of the underlying immune defect.

15 The tat protein, the product of one of the major regulatory genes of the AIDS virus, has been found to be a growth factor for cultured cells derived from Kaposi's sarcoma lesions of AIDS patients. These cells, called spindle cells (KS cells or KS spindle cells) are the
20 suspected tumor cells of KS.

Salahuddin et al., *Science*, 242:430-433 (1988) disclose that AIDS-associated KS and possibly other types of KS may be initiated by signals that induce the growth of these KS cells. AIDS-KS cells cultured in the presence of
25 conditioned medium from HTLV-II-infected and transformed T cell lines were studied. Growth stimulation was induced in the AIDS-KS cells, and not in control cells, suggesting that these KS spindle cells might play an important role in the development and maintenance of KS lesions and, more
30 importantly, that a factor released by HTLV-infected and transformed T cell lines was responsible for stimulating the AIDS-KS cells.

Ensoli et al., *Nature*, 345:84-86 (1990) disclose that tat is released into the medium from both HIV-1-
35 acutely infected H9 cells and COS-1 cells transfected with the tat gene. tat-containing medium specifically promoted

-22-

AIDS-KS cells (cultured spindle-like cells derived from KS lesions of AIDS patients) which were inhibited by anti-tat antibodies indicating that extracellular tat could be involved in the development or progression, or both, of KS in HIV-1-infected individuals. Transplantation of AIDS-KS cells into nude mice produced mouse lesions closely resembling KS.

The presence of KS-growth promoting activity in conditioned media from HIV-1-infected CD4+T cells, the absence of HIV-1 sequences in DNA from KS tissue or cultured cells, and the observation that transgenic mice carrying the tat gene develop KS-like lesions and express tat in the skin but not in the tumor cells, indicate that the role of HIV-1 in KS is indirect, and that tat itself might be released by infected cells and promote activation and growth of target cells involved in the formation of KS.

Further evidence for the role of tat in KS has resulted from experiments with transgenic mice. When the tat gene is introduced into mice, the gene is expressed in the skin only. The tat gene expression in the skin of transgenic mice is correlated with the development of skin tumors that closely resemble KS in humans.

Vogel et al., Nature, 335:606-611 (1988) disclose introducing the tat gene under the control of the HIV LTR into the germline of mice. The resulting transgenic animals developed dermal lesions resembling Kaposi's sarcoma (KS) suggesting that HIV, and specifically the tat gene product, contributes to the development of KS.

tat AND NEUROTOXICITY

Infection with HIV-1 is often complicated by neurological syndromes that include dementia, subacute encephalitis, and vacuolar degeneration of the spinal cord. The identification and isolation of HIV-1 from the brain suggests that the retroviral infection is responsible for the neurological disorders observed in HIV-infected patients.

-23-

Sabatier et al., *J. Virol.*, 65:961-967 (1991) disclose that the intracerebroventricular injection of tat or some tat fragments caused neurotoxic and lethal effects in mice. tat neurotoxicity was also investigated by
5 structure-activity relationships, using binding experiments and electrophysiology. The tat binding site is identified as that region from 48 to 66 containing a highly basic domain critical for efficient tat trans-activation. It is shown that tat binds to the membrane-lipid bilayer of cell
10 membrane by its basic domain. It is suggested that tat binding can directly provoke some biological effects such as neural stimulation, promoting neurological dysfunction.
tat AND IMMUNODEFICIENCY

One of the hallmarks of AIDS is depletion of T4
15 cells, with the subsequent development of immunodeficiency. However, destruction of CD4+ T-cells does not adequately explain the immunopathogenic effects of HIV infection. For example, even early in infection, patient lymphocytes have a defect in their ability to recognize and respond to
20 soluble antigens *in vitro*, even though there are still normal numbers of CD4+ T lymphocytes. In contrast, ability of lymphocytes to proliferate in response to mitogens is not lost in these patients. Viscidi et al., *Science*, 246:1606-1608 (1989) disclose that tat inhibits antigen-
25 induced, but not mitogen-induced, lymphocyte proliferation. In *in vitro* studies, 50 nM Tat was sufficient for 50% inhibition, suggesting that Tat may be a potent immunosuppressive agent. Viscidi et al. did not know whether tat must be provided extracellularly or whether tat
30 produced internally can elicit these effects. In the present invention, topical application is believed to be most useful for treatment of KS. However, the form of administration will be dependent on the therapeutic utility.

-24-

Thus it has been shown that:

1. the tat protein and its interaction with TAR RNA is crucial for HIV replication,
2. tat protein secreted from cells appears to play a role in the development or progression of Kaposi's sarcoma in AIDS,
3. the tat protein has specific neurotoxic effects which suggest that binding of tat to membranes of the central nervous system may cause the neurological syndromes often associated with AIDS, and
4. the tat protein inhibits T-cell proliferation in a specific manner which indicates that tat might directly contribute to the immunosuppression associated with AIDS.

15 THE HIV CAR ELEMENT AND THE rev PROTEIN

The HIV CAR element provides another preferred embodiment of the invention. One of the regulatory events in the life cycle of the human immunodeficiency virus is accumulation of the large virion structural RNA's which are accumulated at the expense of the shorter, regulatory RNA's. In essence, the virus uses much of the same RNA material to encode each set of proteins. If the RNA's are more extensively spliced, the regulatory proteins are produced. If the RNA's are less extensively spliced, the structural proteins are produced. For example, See: W.A. Haseltine, F. Wong-Staal, *Scientific American*, October, 52 (1988). These events are regulated by a protein known as rev, which is produced by the rev gene. Rev's function is to enhance the transport of RNA from the nucleus of the cell to the cytoplasm. In the absence of rev, the mRNA's stay in the nucleus of the cell, where they are subject to splicing enzymes which convert them to mRNA's which encode regulatory proteins. In the presence of rev, the mRNA's are transported to the cytoplasm leading to less splicing.

35 Rev functions by binding to an RNA structural

-25-

element known as the CAR element as reported by E.T. Dayton, D.M. Powell, A.I. Dayton, in *Science* 246, 1625 (1989). This structural element has also been referred to as the rre (rev-responsive element). The functional RNA

5 has been localized to a 269 bp region in the env RNA with the coordinates 7358-7627. For example, See: L. Ratner, W. Haseltine, R. Patarca, K.J. Livak, B. Starcich, S.F. Josephs, *Nature* 313, 277 (1985). The linear CAR sequence is shown in Figure 2. For convenience, this structure is

10 referred to as the CAR element. The secondary structure of the CAR element is currently not known with certainty. However, it is possible to predict the secondary structure of the CAR element using computer programs commonly used by those skilled in the art such as the program of Zuker as

15 described in M. Zuker, *Science*, 244, 48 (1989). The result of such an analysis yielded the result shown in Figure 3. Each of the stem loop structures shown in Figure 3 has the potential to interact with the rev gene product and each can be mimicked by oligonucleotides or oligonucleotide

20 analogs as an embodiment of this invention. It is by no means certain that the structures predicted by the computer program and illustrated in Figure 3 are correct. This does not restrict the practice of this invention for the CAR element structure, however. In this and all other cases

25 where the actual RNA structure is uncertain, the invention can be practiced by preparing a series of oligonucleotides or oligonucleotide analogs which scan the sequence, beginning with the structures predicted to have the lowest energy according to the computer predictions, and proceed

30 to make additional oligonucleotide or oligonucleotide compositions sequentially to the less energetically favored structures.

Assays to measure the normal function of the rev gene product can be conveniently performed according to

35 published procedures. See Dayton et al., J. Acq. Immune Deficiency Syndromes, 1, 441, 1988. Briefly, vectors which

-26-

express HIV mRNA in cells under regulatory control of a variety of promoters are transfected into cells along with a vector which expresses the rev protein. When rev functions normally to facilitate the transport of mRNA to the cytoplasm, the transported mRNA's encode the gag protein, which is detected by an immunoabsorbant assay. When oligonucleotides or oligonucleotide analogs interfere with this process, a decrease in production of gag protein is measured. The reagents needed to conduct these experiments are available from the National Institutes of Health through the AIDS Research and Reference Reagent Program, 1990 catalog, National Institute of Allergy and Infectious Diseases.

The effects of oligonucleotides and oligonucleotide analogs will be determined by adding the compounds directly to the transfection mixture or by adding the compounds to the media at various times and concentrations following transfection, followed by the assay at, for example, 24-48 hours post-transfection.

The present invention relates to compounds which are believed to specifically mimic HIV RNA structures and interfere with viral replication and function. These oligonucleotide and oligonucleotide analog compounds have been shown to have activity in modulating the expression of certain HIV proteins. In accordance with the teachings of the invention, the following examples are provided relating to oligonucleotide synthesis, purification and analysis, including specific oligonucleotide sequences and configurations; and cell-based evaluations of these exemplary oligonucleotides.

EXAMPLES

EXAMPLE 1

OLIGONUCLEOTIDE SYNTHESIS AND PURIFICATION

Synthesis

Unmodified oligonucleotides were synthesized on an Applied Biosystems 380B DNA Synthesizer using standard

-27-

phosphoramidite chemistry with oxidation by iodine. The reagents, both CPG-bound and β -cyanoethyl-diisopropyl-phosphites, were purchased from Applied Biosystems, Inc. (Foster City, CA). For preparation of phosphorothioate oligonucleotides, the standard oxidation bottle was replaced by 0.2 M solution of 3 H-1,2-benzodithiole-3-one 1,1-dioxide (R.P. Iyer, W. Egan, J.B. Regan, and S.L. Beaucage, *J. Am. Chem. Soc.* (1990) 112:1253-1254) in acetonitrile for the stepwise thiation of the phosphoramidite linkages. The thiation cycle time was increased to 68 seconds. After cleavage from the CPG-column and deblocking in concentrated ammonium hydroxide at 55°C (18 hours), the phosphorothioates were purified by trityl-on HPLC with a PRP-1 column using a gradient of acetonitrile in 50 mM of triethyl-ammonium acetate, pH 7 (4% to 32% in 30 minutes, flow rate of 1.5 ml/minute). Appropriate fractions were pooled, evaporated, and treated with 5% acetic acid at ambient temperature for 15 minutes. The solution was extracted with an equal volume of ethyl acetate, neutralized with ammonium hydroxide, frozen, and lyophilized. For the preparation of 2'-O-Me oligonucleotides, the normal phosphoramidite monomers were replaced with 2'-O-Me-substituted phosphoramidites purchased from Chemgenes. Analytical gel electrophoresis was accomplished in 20% AA, 8 M urea, 45 mM trisborate buffer, pH 7, 40 V/cm.

NMR Analysis of Oligonucleotides

The relative amounts of phosphorothioate and phosphodiester linkages obtained by our synthesis were determined by ^{31}P NMR spectroscopy. The spectra were acquired on a Varian NMR spectrometer with a ^{31}P frequency of 162 MHz. Typically, 1,000 transients are coadded. A relaxation delay of 7.5 seconds between transients is used to insure a fully relaxed spectrum. The ^{31}P spectra are acquired at ambient temperature using deuterium oxide or dimethyl sulfoxide- d_6 as a solvent. Phosphorothioate

-28-

samples typically contained less than one percent of phosphordiester linkages.

EXAMPLE 2

MANAGEMENT OF HIV LTR GENE EXPRESSION IN CELLS

5 HeLa cells were maintained in DMEM plus 10% FCS. For antisense experiments, cells were seeded in 6 well dishes at 50% confluency the day prior to the experiment. For each dish, 1 μ g of pHIVpap and 12 μ g of pcDEBtat were precipitated in 500 μ l of 1X HBS and 32 μ l of 2.5 M CaCl_2 .
10 The CaPO_4 precipitate was then evenly divided between the six wells and allowed to sit for 6 hours prior to removal of the precipitate and addition of fresh media. Sixteen hours later, antisense oligonucleotides were transfected by the same method. The next day cells were washed 2X with
15 TBS and harvested in 500 μ l of which 100 μ l were used in the protein assay. The remaining 400 μ l of cell suspension was pelleted, then resuspended in 50 μ l TBS buffer. Next, endogenous phosphatases were inactivated by heating to 65°C for 30 minutes. The heat stable human placental alkaline
20 phosphatase activity was then assayed by the addition of 500 μ l of 5mM PNPP (Sigma) in TEA buffer to the cell suspension, followed by incubation at 37°C. Phosphatase activity was determined at 30 minute intervals using 150 μ l aliquots of the reaction mixture and measuring the
25 absorbance at 405 nm with a Titertek Multiscan MCC/340 ELISA plate reader. The PAP activity was normalized to total protein in each well as determined by Bio-Rad protein assay, in which 1/5 of the harvested cells in TBS (0.1 μ l) were added to 30 μ l of Bio-Rad protein reagent, then
30 incubated for 10 minutes at room temperature, followed by measurement of absorbance at 595 nm using the Titertek plate reader. The inhibition of PAP activity observed with oligonucleotides 1345, 1346, 1347, 1348 and 1349 are shown in Figures 5 and 6.

EXAMPLE 3**TAR RNA SYNTHESIS**

Template synthesis. A duplex DNA template for T7 RNA polymerase copying into TAR RNA was synthesized using the polymerase chain reaction (PCR). PCR primers were designed to be complementary to sequences of pHIV-PAP: the 5'-primer (35-mer) construction consisted of 5'-most sequences corresponding to the 17 base T7 RNA polymerase promoter, and were non-homologous to pHIV-PAP sequences, followed by 18 bases which were homologous to the first 18 nucleotides of the TAR stem-loop (5'-AAT AGC ACT CAC TAT AGG GTC TCT CTG GTT AGA CCA-3'); the 3'-primer (20-mer) was homologous to the last 20 bases of the TAR stem-loop (5'-CCA GCA TGT CTG GAG GGC AG-3'). A standard PCR setup with Taq polymerase and 30 cycles of amplification was used. Amplified duplex template was purified by standard organic solvent extractions followed by ethanol/sodium acetate precipitation.

T7 RNA polymerase synthesis of TAR RNA. Duplex DNA template was added at 500pM to a 1.0 mL reaction mix containing 1 2.5mM each of GTP, CTP, ATP & UTP, 40mM tris-HCl pH 8.0, 1 .0mM spermidine, 5mM DTT, 0.01% (v/v) triton X-100, 20% (v/v) PEG 8000, 31 mM MgCl₂, and 10% (v/v addition previously optimized for polymerase reaction efficiency) of a T7 RNA polymerase preparation. The reaction was incubated for 4h at 37°C. The RNA product was PAGE purified, dephosphorylated with calf intestinal alkaline phosphatase, concentration determined by UV absorbance, and 5'-end-labeled with ³²P to high specific radioactivity (7000Ci/mmol) using T4 polynucleotide kinase.

EXAMPLE 4**POLYACRYLAMIDE GEL MOBILITY SHIFT ASSAYS OF TAR-tat BINDING**

All dilutions of [5'-³²P]-TAR RNA stock solution were made into TE, pH 7.5 containing 4mg pdldC, and all dilutions of stock solutions (in water) of Tat 39-mer

-30-

peptide (Tat3B; from 48-86 in the Tat protein sequence) similarly were made into TE, pH 7.5 containing 500nM BSA. Inclusion of pdldC and BSA was made to reduce the effect of nonspecific adsorption (to solid surfaces) of dilute solutions of RNA and peptide, respectively, and independently were shown to be without effect on the specific binding of Tat39 to TAR RNA. Tat39 used was obtained from the UCSF Biotechnology Resource Core facility and was made by solid phase automated chemical synthesis, purified by RP HPLC, and characterized by amino acid analysis and mass spectrometry. TAR and analogs not made by T7 RNA polymerase synthesis were made by automated chemical oligonucleotide synthesis (indicated by synthesis number in Table 1). Gel mobility shift assays were performed by the addition of [5'-³²P]-TAR RNA and Tat39 at indicated concentrations (Table 1) to a 10 μ l reaction containing 10mM tris-HCl pH 7.5, 70mM NaCl, 0.2mM EDTA, 5% (v/v) glycerol, 500nM BSA, and 40mg pdldC. Each binding mix was incubated for 30 min at 4°C and then loaded directly onto a 10% (75:1 acrylamide:bisacrylamide) native PAG. Electrophoresis was performed using 1/2 TBE running buffer and 250V at 43°C for ca. 2h. Radiolabeled TAR RNA was then detected by autoradiography. For preliminary screening experiments, binding parameters were determined by visual estimation of relative intensities of exposure of film by free and specifically bound TAR and TAR mimetics/analog, followed by appropriate mathematical and graphical analysis. The results are shown in Table 2.

-31-

TABLE 2
GEL MOBILITY-SHIFT tat-"TAR" BINDING ASSAYS

5	OLIGO	[³² P-OLIGO][Tat39]	1:1KD (nm)	REL (TAR 1:1 K _D) (OLIGO 1:1 K _D)	2-3:1 KD (nm)	SITES (2:1 K _D /1:1 K _D)
10	TAR 59-mer 1-59	5-5000	10-500	40	---	200 5
15	U-DNA TAR #1973 40-mer 11-50	5	10-1000	ND	<0.001	ND
20	A:P=S TAR 59-mer 1-59	5	10-1000	<10	4	30 3
20	loopless TAR #2002/#2246 23-mer 16-29(-A17) +36-45	5*/5000 no duplex 5000*/5x10 ⁶ duplex	10-1000	ND	<0.001	ND
25	5-BrU TAR 59-mer 1-59	5	10-1000	40	1	ND

-32-

5	2'-OMe #2306 29-mer 16-45(-A17)	TAR	5	10-1000	10	4	100	10
	2'-OMe #2306 29-mer 16-45(-A17)	TAR 06 (aged)	50	0.125- 16.4x10 ³	125	0.32	8.2x10 ³	66
10	2'-OMe, P=S #2195 29-mer 16-45(-A17)	TAR	5	10-1000	<10	4	ND	>100
15	2'-OMe, P=S #2195 29-mer 16-45(-A17)	TAR	50	0.125- 16.4x10 ³	4	10	8.0x10 ³ 20x10 ³	2000

* When duplex structures are formed by intermolecular association and hybridization of individual oligos, rather than by intramolecular hybridization of a single oligo, then only one of the oligos is 5"-end-labeled with ³²P and is distinguished from the unlabeled oligo by an .

-33-

EXAMPLE 5**DEVELOPMENT OF LUCIFERASE ASSAY FOR HIV GENE EXPRESSION**

pHIVluc is a plasmid which contains the luciferase gene under regulatory control of the HIV LTR.

5 When this plasmid is present in the cell, it responds to the same regulatory signals which activate HIV gene expression by producing the enzyme luciferase. Luciferase can be easily assayed by adding a substrate, luciferin, under the appropriate conditions and measuring the amount

10 of light produced in a luminometer. Thus, specific inhibition of luciferase production in cells is equivalent to inhibiting HIV gene expression, and is predictive of antiviral activity in humans. This assay is similar to that described by Felber B.K. and Pavlakis G.N. (*Science*,

15 239:184-187 (1988)), the major difference is simply the enzyme encoded by the reporter gene. To construct this plasmid, the plasmids pT3/T7luc (Clonetech) and IP-RG-24 (a plasmid which contains the HIV LTR) were digested to completion with KpnI and HindIII. Restriction fragments

20 containing the luciferase cDNA and the HIV LTR and other processing signals were isolated and ligated to generate pHIVluc, which expresses the luciferase protein under the control of the HIV LTR.

EXAMPLE 6

25 **PROCEDURE FOR MEASURING INHIBITION OF HIV GENE
EXPRESSION IN CULTURED CELLS**

To test for inhibition of HIV gene expression, HeLa cells were seeded at 3×10^5 cells per well of a 6 well plate 16 h prior to the experiment. Test compounds were

30 added to triplicate wells at indicated concentrations. Following a 3 hour incubation the cells were calcium phosphate transfected with pHIVluc and pcDEBtat (Feng S. and Holland E.C., *Nature*, 334:165-167 (1988), which expresses tat, the HIV trans-activator protein. Briefly, 5

-34-

ug of pHIVluc and 6 ug of pcDEBtat were added to 500 ul of 250 mM CaCl_2 , then 500 ul of 2 x HBS were added followed by vortexing. After 30 minutes the DNA precipitate was divided evenly between the six wells of the plate, which was then incubated for 4-6 hours. The media and precipitate were then removed, the cells washed with PBS, and fresh media containing the test compound at the initial concentration was added and incubated for 16 hours.

Luciferase activity was then determined for each well as follows. Media was removed, then the cells washed 2X with PBS. The cells were then lysed on the plate with 200 ul of LB (1% Triton X-100, 25 mM glycylglycine pH 7.8, 15 mM MgSO_4 , 4 mM EGTA, 1mM DTT). A 75 ul aliquot from each well was added to 96 well plate along with 75 ul of assay buffer (25mM glycylglycine pH 7.8, 15 mM MgSO_4 , 4 mM EGTA, 15 mM KPO_4 , 1 mM DTT, 2.5 mM ATP). The plate was then read in a Dynatec multiwell luminometer which injected 75 ul of Luciferin buffer (25 mM glycylglycine pH 7.8, 15 mM MgSO_4 , 4mM EGTA, 4 DTT, 1 mM luciferin) into each well and measured the light emitted.

EXAMPLE 7

ACTIVITY OF RNA MIMETICS IN INHIBITION OF HIV GENE EXPRESSION

Compound 2306 is a 2'-O-methyl oligonucleotide analog 29-mer which forms a truncated HIV TAR stem/loop structure. It was found that compound 2306 binds to the tat peptide *in vitro*. Compound 2848 and 2850 are also 2'-O-methyl analogs of similar length which form stem/loop structures, but are unable to bind tat peptide *in vitro* due to extensive mutations in the loop and bulge regions. In the HIV gene expression assay, compound 2306 shows significant activity in inhibition of HIV gene expression over the controls 2848 and 2850 compounds at doses below $1\mu\text{M}$ (Fig. 7). At a higher dose ($7\mu\text{M}$) there was some non-

-35-

specific activity in the control compounds 2848 and 2850,
which was less than the specific compound 2306.

-36-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Ecker et al.
- 5 (ii) TITLE OF INVENTION: REAGENTS AND METHODS FOR MODULATING
GENE EXPRESSION THROUGH RNA MIMICRY
- (iii) NUMBER OF SEQUENCES: 5
- (iv) CORRESPONDENCE ADDRESS:
- (A) ADDRESSEE: Woodcock Washburn Kurtz
Mackiewicz & Norris
- 10 (B) STREET: One Liberty Place - 46th Floor
- (C) CITY: Philadelphia
- (D) STATE: PA
- (E) COUNTRY: USA
- (F) ZIP: 19103
- 15 (v) COMPUTER READABLE FORM:
- (A) MEDIUM TYPE: DISKETTE, 3.5 INCH, 1.44 Mb STORAGE
- (B) COMPUTER: IBM PS/2
- (C) OPERATING SYSTEM: PC-DOS
- (D) SOFTWARE: WORDPERFECT 5.0
- 20 (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER: n/a
- (B) FILING DATE: herewith
- (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
- 25 (A) APPLICATION NUMBER: 497,090
- (B) FILING DATE: March 21, 1990

-37-

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Jane Massey Licata

(B) REGISTRATION NUMBER: 32,257

(C) REFERENCE/DOCKET NUMBER: ISIS-0109

5 (ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (215) 568-3100

(B) TELEFAX: (215) 568-3439

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 59

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

15 GGGUCUCUCU GGUUAGACCA GAUCUGAGCC UGGGAGCUCU CUGGCUAACU 50
AGGGAACCC

9

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 40

20 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

GGUUAGACCA GAUCUGAGCC UGGGAGCUCU CUGGCUAACU

40

-38-

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13

(B) TYPE: nucleic acid

5 (C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GCCAGAUCUG AGC

13

(2) INFORMATION FOR SEQ ID NO: 4:

10 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

GCUCUCUGGC

10

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29

20 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

GCCAGAUCUG AGCCUGGGAG CUCUCUGGC

29

WHAT IS CLAIMED IS:

1. A method for modulating expression of a gene comprising:
selecting a portion of RNA coded by the gene, said RNA being
capable of interacting with protein;
5 preparing an oligonucleotide or oligonucleotide analog to
mimic said portion; and
contacting cells containing the gene with said oligonucleotide
or oligonucleotide analog.
2. The method of Claim 1 wherein said gene is of an
10 infectious organism.
3. The method of Claim 2 wherein said protein is produced by
a second portion of RNA coded by the infectious organism.
4. The method of Claim 1 wherein interaction of the protein
and the RNA portion effects stimulation of expression of the gene.
- 15 5. The method of Claim 1 wherein said oligonucleotide or
oligonucleotide analog mimics at least about 6 nucleotide units.
6. The method of Claim 1 wherein said oligonucleotide or
oligonucleotide analog mimics from about 8 to about 50 nucleotide
units.
- 20 7. The method of Claim 1 wherein said oligonucleotide or
oligonucleotide analog mimics from about 10 to about 20 nucleotide
units.
8. The method of Claim 1 wherein said RNA possesses a
secondary structure.

9. The method of Claim 8 wherein said oligonucleotide or oligonucleotide analog reproduces at least a portion of said secondary structure.

10. A method for treating a disease comprising:

5 selecting a portion of RNA coded by a gene whose expression is believed to be responsible for said disease, said RNA being capable of interacting with protein;

preparing an oligonucleotide or oligonucleotide analog to mimic said portion; and

10 contacting an organism suspected of having the disease with said oligonucleotide or oligonucleotide analog.

11. The method of Claim 10 wherein said gene is of an infectious organism.

12. The method of Claim 11 wherein said protein is produced
15 by a second portion of RNA coded by the infectious organism.

13. The method of Claim 10 wherein interaction of the protein and the RNA portion effects stimulation of expression of the gene.

14. The method of Claim 10 wherein said oligonucleotide or oligonucleotide analog mimics at least about 6 nucleotide units.

20 15. The method of Claim 10 wherein said oligonucleotide or oligonucleotide analog mimics from about 8 to about 50 nucleotide units.

16. The method of Claim 10 wherein said oligonucleotide or oligonucleotide analog mimics from about 10 to about 20 nucleotide
25 units.

17. The method of Claim 10 wherein said RNA possesses a secondary structure.

-41-

18. The method of Claim 17 wherein said oligonucleotide or oligonucleotide analog reproduces at least a portion of said secondary structure.

19. The method of Claim 10 wherein said contacting is in amounts and for times effective to modulate expression of said gene.

20. The method of Claim 10 wherein said disease is human immunodeficiency virus infection.

21. An oligonucleotide or oligonucleotide analog comprising the sequence:

5'- CU GGG A -3'.

22. The oligonucleotide or oligonucleotide analog of claim 21 further comprising, immediately 5' to said sequence, at least a 3' portion of the sequence:

15 5'- UCU GAG C -3'.

23. The oligonucleotide or oligonucleotide analog of claim 21 further comprising, immediately 3' to said sequence, at least a 5' portion of the sequence:

5'- GC UC -3'.

20 24. The oligonucleotide or oligonucleotide analog of Claim 21 having at least one of the 5' and 3' termini capped to inhibit depolymerization or destruction of the oligonucleotide or oligonucleotide analog.

25. The oligonucleotide or oligonucleotide analog of Claim 21 comprising oligonucleotide analog subunits having improved nuclease resistance as compared with naturally occurring subunits.

-42-

26. The oligonucleotide or oligonucleotide analog of Claim 21 in a pharmaceutically acceptable carrier.

27. An oligonucleotide or oligonucleotide analog comprising the sequence:

5 5'- UCU GAG CCU GGG AGC UC -3'.

28. The oligonucleotide or oligonucleotide analog of Claim 27 further comprising, immediately 5' to the original sequence, at least a 3' portion of the sequence:

5'- CCAGA -3'.

10 29. The oligonucleotide or oligonucleotide analog of Claim 27 further comprising, immediately 3' to the original sequence, at least a 5' portion of the sequence:

5'- GGUCU -3'.

30. The oligonucleotide or oligonucleotide analog of Claim 27 having at least one of the 5' and 3' termini capped to inhibit depolymerization or destruction of the oligonucleotide or oligonucleotide analog.

31. The oligonucleotide or oligonucleotide analog of Claim 27 comprising oligonucleotide analog subunits having improved nuclease resistance as compared with naturally occurring subunits.

32. The oligonucleotide or oligonucleotide analog of Claim 27 in a pharmaceutically acceptable carrier.

33. An oligonucleotide or oligonucleotide analog comprising the sequence:

25 5'- UCU GAG CCU GGG AGC UCA GA -3'.

-43-

34. A method for interfering with the function or replication of a retrovirus comprising contacting said virus or a medium or tissue in which it resides with an oligonucleotide or oligonucleotide analog comprising the sequence:

5 5'- CU GGG A -3'.

35. The method of Claim 34 wherein said oligonucleotide or oligonucleotide analog further comprises, immediately 5' to the original sequence, at least a 3' portion of the sequence:

5'- UCU GAG C -3'.

10 36. The method of Claim 34 wherein said oligonucleotide or oligonucleotide analog further comprises, immediately 3' to the original sequence, at least a 5' portion of the sequence:

5'- GC UC -3'.

37. The method of Claim 34 wherein said oligonucleotide or oligonucleotide analog comprises oligonucleotide analog subunits having improved nuclease resistance as compared with naturally occurring subunits.

38. The method of Claim 34 wherein said oligonucleotide or oligonucleotide analog is in a pharmaceutically acceptable carrier.

20 39. The method of Claim 34 wherein said retrovirus is a human immunodeficiency virus.

40. A method for interfering with the function or replication of a retrovirus comprising contacting said virus or a medium or tissue in which it resides with an oligonucleotide or oligonucleotide analog comprising the sequence:

5'- UCU GAG CCU GGG AGC UC -3'.

-44-

41. The method of Claim 40 wherein said oligonucleotide or oligonucleotide analog further comprises, immediately 5' to the original sequence, at least a 3' portion of the sequence:

5'- CCAGA -3'.

5 42. The method of Claim 40 wherein said oligonucleotide or oligonucleotide analog further comprises, immediately 3' to the original sequence, at least a 5' portion of the sequence:

5'- GGUCU -3'.

43. The method of Claim 40 wherein said oligonucleotide or oligonucleotide analog comprises oligonucleotide analog subunits having improved nuclease resistance as compared with naturally occurring subunits.

44. The method of Claim 40 wherein said oligonucleotide or oligonucleotide analog is in a pharmaceutically acceptable carrier.

15 45. The method of Claim 40 wherein said retrovirus is a human immunodeficiency virus.

46. A method for treating an animal suspected of being infected with an immunodeficiency virus comprising contacting the animal with an oligonucleotide or oligonucleotide analog comprising
20 the sequence:

5'- CU GGG A -3'.

47. The method of Claim 46 wherein said oligonucleotide or oligonucleotide analog further comprises, immediately 5' to the original sequence, at least a 3' portion of the sequence:

-45-

5'- UCU GAG C -3'.

48. The method of Claim 46 wherein said oligonucleotide or oligonucleotide analog further comprises, immediately 3' to the original sequence, at least a 5' portion of the sequence:

5 5'- GC UC -3'.

49. The method of Claim 46 wherein said oligonucleotide or oligonucleotide analog comprises oligonucleotide analog subunits having improved nuclease resistance as compared with naturally occurring subunits.

10 50. The method of Claim 46 wherein said oligonucleotide or oligonucleotide analog is in a pharmaceutically acceptable carrier.

51. A method for treating an animal suspected of being infected with an immunodeficiency virus comprising contacting the animal with an oligonucleotide or oligonucleotide analog comprising
15 the sequence:

5'- UCU GAG CCU GGG AGC UC -3'.

52. The method of Claim 51 wherein said oligonucleotide or oligonucleotide analog further comprises, immediately 5' to the original sequence, at least a 3' portion of the sequence:

20 5'- CCAGA -3'.

53. The method of Claim 51 wherein said oligonucleotide or oligonucleotide analog further comprises, immediately 3' to the original sequence, at least a 5' portion of the sequence:

5'- GGUCU -3'.

-46-

54. The method of Claim 51 wherein said oligonucleotide or oligonucleotide analog comprises oligonucleotide analog subunits having improved nuclease resistance as compared with naturally occurring subunits.

5 55. The method of Claim 51 wherein said oligonucleotide or oligonucleotide analog is in a pharmaceutically acceptable carrier.

56. A method for treating an animal suspected of being infected with an immunodeficiency virus comprising contacting the animal with an oligonucleotide or oligonucleotide analog comprising the sequence:

5'- UCU GAG CCU GGG AGC UCA GA -3'.

57. A method for treating human immunodeficiency virus infection comprising administering to a patient suspected of having said infection an oligonucleotide or oligonucleotide analog mimic of at least a portion of the TAR region of the HIV mRNA.

58. A method for treating human immunodeficiency virus infection comprising administering to a patient suspected of having said infection an oligonucleotide or oligonucleotide analog mimic of at least a portion of the CAR region of the HIV mRNA.

20 59. A method for treating human immunodeficiency virus infection comprising administering to a patient suspected of having said infection an oligonucleotide or oligonucleotide analog mimic of at least a portion of the gag-pol region of the HIV mRNA.

60. Oligonucleotide or oligonucleotide analog comprising one of the sequences :

5'-GGG UCU CUC UGG UUA GAC CAG

AUC UGA GCC UGG GAG CUC UCU

-47-

GGC UAA CUA GGG AAC CC-3', SEQ. ID NO: 1;

5'-GGU UAG ACC AGA UCU GAG CCU

GGG AGC UCU CUG GCU AAC U-3', SEQ. ID NO: 2;

5'- GCC AGA UCU GAG C-3', SEQ. ID NO: 3;

5 5'-GCU CUC UGG C-3', SEQ. ID NO: 4; and

5'-GCC AGA UCU GAG CCU GGG AGC

UCU CUG GC-3', SEQ. ID NO: 5.

61. Oligonucleotide or oligonucleotide analog of Claim 60 in a pharmaceutically acceptable carrier.

10 62. Method for interfering with the function or replication of a retrovirus comprising contacting said virus or a medium or tissue in which said virus resides with an oligonucleotide or oligonucleotide analog comprising one of the sequences:

5'-GGG UCU CUC UGG UUA GAC CAG

15 AUC UGA GCC UGG GAG CUC UCU

GGC UAA CUA GGG AAC CC-3', SEQ. ID NO: 1;

5'-GGU UAG ACC AGA UCU GAG CCU

GGG AGC UCU CUG GCU AAC U-3', SEQ. ID NO: 2;

-48-

5'- GCC AGA UCU GAG C-3', SEQ. ID NO: 3;

5'-GCU CUC UGG C-3', SEQ. ID NO: 4; and

5'-GCC AGA UCU GAG CCU GGG AGC

UCU CUG GC-3', SEQ. ID NO: 5.

5 63. Method of Claim 62 wherein said oligonucleotide or oligonucleotide analog is in a pharmaceutically acceptable carrier.

64. Method for treating an animal suspected of being infected with an immunodeficiency virus comprising contacting the animal with an oligonucleotide or oligonucleotide analog comprising one of the sequences:

5'-GGG UCU CUC UGG UUA GAC CAG

AUC UGA GCC UGG GAG CUC UCU

GGC UAA CUA GGG AAC CC-3', SEQ. ID NO: 1;

5'-GGU UAG ACC AGA UCU GAG CCU

15 GGG AGC UCU CUG GCU AAC U-3', SEQ. ID NO: 2;

5'- GCC AGA UCU GAG C-3', SEQ. ID NO: 3;

5'-GCU CUC UGG C-3', SEQ. ID NO: 4; and

5'-GCC AGA UCU GAG CCU GGG AGC

UCU CUG GC-3', SEQ. ID NO: 5.

65. Method of Claim 64 wherein said oligonucleotide or oligonucleotide analog is in a pharmaceutically acceptable carrier.

FIGURE 1A

5' 3'
 GGGUCUCUCUGGUUAGACCAGAUCUGAGCCUGGGAGCUCUCUGGCUAACUAGGGAACCC

FIGURE 1B

	C		A	UCU	CU
5'	-GGGU	UCUCUGGUUAG	CCAGA	GAGC	G
3'	-CCCA	AGGGAUCAAUC	GGUCU	CUCG	G
	-	-	---	AG	

2/7

FIGURE 2

[7357]

UCCUUGGGUU CUUGGGAGCA GCAGGAAGCA CUAUGGGCGC AGCGUCAAUG

ACGCUGACGG UACAGGCCAG ACAAUUAUUG UCUGGUUAUAG UGCAGCAGCA

GAACAAUUG CUGAGGGCUA UUGAGGCGCA ACAGCAUCUG UUGCAACUCA


CAGUCUGGGG CAUCAAGCAG CUCCAGGCAA GAAUCCUGGC UGUGGAAAGA

UACCUIAAAGG AUCAACAGCU CCUAGGGAUU UGGGGUUGCU CUGGAAAACU

CAUUUGCACC ACUGCUGUGC

[7627]

3/7

***Fig. 3***

SUBSTITUTE SHEET

4/7





Fig. 4

5/7

**Fig. 5**

6/7



**Fig. 6**

FIGURE 7



6 ug pHIVluc, 4 ug pcDEBlat

INTERNATIONAL SEARCH REPORT

International Application No. **PCT/US91/01822**

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC IPC(5): A61K 31/70; C07H 15/12, 17/00; C12N 7/04 U.S. Cl.: 514/44; 435/236; 536/27		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
U. S.	514/44; 435/236; 536/27	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
DIALOG DATABASES: BIOSIS PREVIEWS 1985+, MEDLINE 1985+		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
P, A	Trends in Genetics, Vol. 7, No. 1, issued January 1991, C. A. Rosen, "Regulation of HIV gene expression by RNA-protein interactions," pages 9-14. See entire article.	1-65
P, Y	Cell, Vol. 63, issued 02 November 1990, B. A. Sullenger et al, "Overexpression of TAR Sequences Renders Cells Resistant to Human Immunodeficiency Virus Replication," pages 601-608. See entire article.	1-57, 60-65
Y	Nature, Vol. 335, issued 29, September 1988, D. Baltimore, "Intracellular Immunization," pages 395-396. See page 395, 3rd column.	1-65
Y	J. Virol., Vol. 63 (No. 3), issued March 1989, M. Hadzopoulou-Cladaras et al, "The <u>rev</u> (<u>tra/art</u>) Protein of Human Immunodeficiency Virus Type 1 Affects Viral mRNA and Protein Expression via a <u>cis</u> -Acting Sequence in the <u>env</u> Region," pages 1265-1274, see page 1272, 2nd column, 3rd full paragraph.	1-56, 58, 60-65
<div style="display: flex; justify-content: space-between;"> <div style="width: 48%;"> <p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 48%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"Z" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
05 JUNE 1991		19 JUN 1991
International Searching Authority		Signature of Authorised Officer
ISA/US		<i>Johnny F. Railey II</i> JOHNNY F. RAILEY II

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category*	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
Y	Proc. Natl. Acad. Sci., Vol. 86, issued September 1989, C. Dingwall et al, "Human immunodeficiency virus 1 tat protein binds trans-activation-response region (TAR) RNA <u>in vitro</u> ," pages 6925-6929. See page 6928, 2nd column, 1st full paragraph and Discussion.	1-57, 60-65

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

- | | | |
|---|--|-----------------|
| Y | Cell, Vol. 60, issued 23 February 1990, M. H. Malim et al, "HIV-1 Structural Gene Expression Requires Binding of the Rev Trans-activator to Its RNA Target Sequence," pages 675-683. See pages 678-680, Figure 4 and Discussion. | 1-56, 58, 60-65 |
| Y | Science, Vol. 247, issued 16 February 1990, H. S. Olsen et al, "Secondary Structure Is the Major Determinant for Interaction of HIV rev Protein with RNA," pages 845-848. See page 847, 3rd column and Figure 3. | 1-56, 58, 60-65 |

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE¹

This International search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers _____, because they relate to subject matter ¹² not required to be searched by this Authority, namely:

2. ☐ Claim numbers _____, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out ¹³, specifically:

3. ☐ Claim numbers _____, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING²

This International Searching Authority found multiple inventions in this international application as follows:

SEE ATTACHMENT

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remarks on Protest

- ☐ The additional search fees were accompanied by applicant's protest
- ☒ No protest accompanied the payment of additional search fees

Attachment to Form PCT/ISA/210, Part VI
Continuation of OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

- Group I: Claims 1-9, 21-33 and 60-61 are drawn to a first product, oligonucleotides or oligonucleotide analogs; and to a method for modulating expression of a gene using oligonucleotides or oligonucleotide analogs.
- Group II: Claims 10-20 are drawn to a second method of use of oligonucleotides or oligonucleotide analogs for treating a disease.
- Group III: Claims 34-45 and 62-63 are drawn to a third method of use of oligonucleotides or oligonucleotide analogs for interfering with the function or replication of a retrovirus.
- Group IV: Claims 46-56 and 64-65 are drawn to a fourth method of use of oligonucleotides or oligonucleotide analogs for treating an animal suspected of being infected with an immunodeficiency virus.
- Group V: Claim 57 is drawn to a fifth method of use of oligonucleotide or oligonucleotide analog mimics of a portion of the TAR region of the HIV mRNA for treating human immunodeficiency virus infection.
- Group VI: Claim 58 is drawn to a sixth method of use of oligonucleotide or oligonucleotide analog mimics of a portion of the CAR region of the HIV mRNA for treating human immunodeficiency virus infection.
- Group VII: Claim 59 is drawn to a seventh method of use of oligonucleotide or oligonucleotide analog mimics of the gag-pol region of the HIV mRNA for treating human immunodeficiency virus infection.

The claims of group I are drawn to a product and a first method of use of the product. The claims of groups II-VII are drawn to distinct methods as described above. PCT Rules 13.1 and 13.2 do not provide for multiple distinct methods within a single general inventive concept. Note also 37 CFR § 1.475.